


CLAIMS

1. A method of treating a disease, damage or disorder of the central nervous system associated with a disorder of neurochemical equilibrium of a biogenic amine or other neurotransmitter, comprising administering to a subject in need thereof a compound of formula **I**




wherein

X is selected from the group consisting of CH₂, O, S, S(=O), S(=O)₂ and NR^a, wherein R^a is selected from the group consisting of hydrogen, C₁-C₃-alkyl, C₁-C₃-alkanoyl, C₁-C₇-alkoxycarbonyl, C₇-C₁₀-arylmethoxycarbonyl, C₇-C₁₀-aroyl, C₇-C₁₀-arylalkyl, C₃-C₇-alkylsilyl and C₅-C₁₀-alkylsilylalkoxyalkyl;

Y and Z are each independently selected from the group consisting of hydrogen, halogen, C₁-C₄-alkyl, C₂-C₄-alkenyl, C₂-C₄-alkynyl, halo-C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkanoyl, amino, amino-C₁-C₄-alkyl, *N*-(C₁-C₄-alkyl)amino, *N,N*-di(C₁-C₄-alkyl)amino, thiol, C₁-C₄-alkylthio, sulfonyl, C₁-C₄-alkylsulfonyl, sulfinyl, C₁-C₄-alkylsulfinyl, carboxy, C₁-C₄-alkoxycarbonyl, cyano and nitro;

R¹ is selected from the group consisting of CHO, CH=CHOCOCH₃, (CH₂)_mOH wherein m is an integer from 1 to 3, and a substituent of the formula **II**:



wherein

R³ and R⁴ are each independently C₁-C₄-alkyl or aryl; or


R³ and R⁴ taken together with the nitrogen atom to which they are attached form a heterocycle or heteroaryl group that is optionally substituted with one or two substituents which are selected from the group consisting of halogen, C₁-C₄ alkyl, cyano,

nitro, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl, and C₁-C₄ alkylsulfinyl;

m is an integer from 1 to 3;

n is an integer from 0 to 3;

Q₁ and Q₂ are each independently selected from the group consisting oxygen, sulfur,



wherein substituents

*y*₁ and *y*₂ are each independently selected from the group consisting of hydrogen, halogen, C₁-C₄-alkyl optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl and C₁-C₄ alkylsulfinyl; hydroxy, C₁-C₄-alkoxy, C₁-C₄-alkanoyl, thiol, C₁-C₄-alkylthio, sulfonyl, C₁-C₄-alkylsulfonyl, sulfinyl, C₁-C₄-alkylsulfinyl, cyano, nitro, and an aryl group wherein said aryl group is optionally substituted with one or two substituents selected from the group consisting of fluoro, chloro, C₁-C₄ alkyl, cyano, nitro, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl, C₁-C₄ alkylsulfinyl and is linked to the rest of the molecule via a direct bond or a C₁-C₄ alkylene group; or

*y*₁ and *y*₂ taken together with the carbon atom to which they are attached form a carbonyl group or an imino group;

R² is selected from the group consisting of hydrogen, a C₁-C₇-alkyl group optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl and C₁-C₄ alkylsulfinyl; ~~or~~ an aryl group optionally substituted with one or two substituents selected from the group consisting of fluoro, chloro, C₁-C₄ alkyl, cyano, nitro, hydroxy, C₁-C₄ alkoxy, thiol, C₁-C₄ alkylthio, amino, *N*-(C₁-C₄) alkylamino, *N,N*-di(C₁-C₄-alkyl)-amino, sulfonyl, C₁-C₄ alkylsulfonyl, sulfinyl, C₁-C₄ alkylsulfinyl; C₁-C₇-alkanoyl, C₁-C₇-alkoxycarbonyl, C₇-C₁₀-

arylalkyloxycarbonyl, C₇-C₁₀-aroyl, C₇-C₁₀-arylalkyl, C₃-C₇-alkylsilyl, C₆H₅CH₂CH₂ and CH₂OCH₂CH₂Si(CH₃)₃;


and of their a pharmaceutically acceptable salt or solvate thereof .

2. The method of claim 1, wherein the biogenic amine is serotonin, norepinephrine or dopamine.
3. The method of claim 1, wherein the neurotransmitter is glutamate.
4. The method of claim 1, wherein the compound of formula **I** regulates the synthesis, storage, release, metabolism, reabsorption, or receptor binding of said biogenic amine or a neurotransmitter.
5. The method of claim 4, wherein the compound of formula **I** binds to a receptor of a biogenic amine.
6. The method of claim 5, wherein the compound of formula **I** binds to a serotonin 5-HT_{2A} or 5-HT_{2C} receptor.
7. The method of claim 6, wherein the compound of formula **I** binds to a serotonin 5-HT_{2A} or 5-HT_{2C} receptor with an IC₅₀ of less than 1 μM.
8. The method of claim 1, wherein the compound of formula **I** binds to a σ1 receptor with an IC₅₀ of less than 1 μM.
9. The method of claim 1, wherein the compound of formula **I** bind to a σ1 receptor and to at least one serotonin receptor selected from 5-HT_{2A} and 5-HT_{2C}.
10. The method of claim 1, wherein the disease or disorder of the central nervous system is selected from the group consisting of anxiety, depression, bipolar disorders, sleeping disorders, sexual disorders, psychosis, borderline psychosis, schizophrenia, migraine, personality disorders, obsessive-compulsive disorders, social phobia, panic attacks, organic mental disorders in children, aggression, memory disorders, personality disorders in elderly people, addiction, obesity, bulimia and other eating disorders, snoring, and premenstrual troubles.
11. The method of claim 1, wherein the damage to the central nervous system is caused by trauma, brain stroke, neurodegenerative diseases, cardiovascular disorders, thrombosis, infarct or gastrointestinal disorders.

12. The method of claim 1 wherein X is O, S, or NR^a, wherein R^a is hydrogen or a substituent selected from the group consisting of C₁-C₃-alkyl, C₁-C₃-alkanoyl, C₇-C₁₀-aroyl, and C₇-C₁₀-arylalkyl.

13. The method of claim 1, wherein Y and Z are each independently selected from the group consisting of hydrogen, fluorine, chlorine, bromine, C₁-C₄-alkyl, halo-C₁-C₄-alkyl, hydroxy, C₁-C₄-alkoxy, trifluoromethoxy, C₁-C₄-alkanoyl, amino, amino-C₁-C₄-alkyl, *N*-(C₁-C₄-alkyl)amino, *N,N*-di(C₁-C₄-alkyl)amino, thiol, C₁-C₄-alkylthio, cyano and nitro.

14. The method of claim 1, wherein R¹ is selected from the group consisting of CHO, CH=CHOCOCH₃, (CH₂)_mOH wherein m is an integer from 1 to 3; ϕ and a substituent of the formula **II**:



II

wherein

R³ and R⁴ are each independently hydrogen, C₁-C₄-alkyl, or aryl or

R³ and R⁴ taken together with the nitrogen atom to which they are attached form a heterocycle or heteroaryl group selected from the group consisting of morpholine-4-yl, piperidine-1-yl, pyrrolidine-1-yl, imidazole-1-yl and piperazine-1-yl;

m is an integer from 1 to 3;

n is an integer from 0 to 3; and

Q₁ and Q₂ are each independently oxygen or CH₂.

15. The method of claim 1, wherein the compound of formula **I**, is selected from the group consisting of:

1-methyl-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

1-methyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

1-phenethyl-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

1-phenethyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

5-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

5-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulene-2-carbaldehyde;

3-(1-phenethyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-yl)-acrylic acid methyl ester;

(1-methyl-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-yl)-methanol;

(1-methyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-yl)-methanol;

(1-phenethyl-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-yl)-methanol;

(1-phenethyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-yl)-methanol;

[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-yl]-methanol;

[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-yl]-methanol;

[5-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-yl]-methanol;

[11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-yl]-methanol;

[5-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-yl]-methanol;

[11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-yl]-methanol;

3-(1-phenethyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-yl)-propane-1-ol;

dimethyl-[2-(1-methyl-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-amine;

dimethyl-[3-(1-methyl-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-amine;

dimethyl-[2-(1-methyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-amine;

dimethyl-[3-(1-methyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-amine;

dimethyl-[2-(1-phenethyl-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-amine;

dimethyl-[3-(1-phenethyl-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-amine;

dimethyl-[2-(1-phenethyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-amine;

dimethyl-[3-(1-phenethyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-amine;

dimethyl-{2-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-ethyl}-amine;

dimethyl-[2-(1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-amine;

dimethyl-{3-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-ylmethoxy]-propyl}-amine;

dimethyl-[3-(1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-amine;

3-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-propylamine;

3-(1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propylamine;

dimethyl-{2-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-ethyl}-amine;

dimethyl-[2-(1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-amine;

dimethyl-{3-[1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-propyl}-amine;

dimethyl-[3-(1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-amine;

{3-[5-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-propyl}-dimethyl-amine;

[3-(5-chloro-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-dimethyl-amine;

3-[5-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-propylamine;

3-(5-chloro-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propylamine;

{2-[11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-ethyl}-dimethyl-amine;

[2-(11-chloro-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-dimethyl-amine;

{3-[11-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-propyl}-dimethyl-amine;

[3-(11-chloro-1H-8-oxa-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-dimethyl-amine;

{2-[5-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-ethyl}-dimethyl-amine;

[2-(5-chloro-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-ethyl]-dimethyl-amine;

{3-[5-chloro-1-(2-trimethylsilyl-ethoxymethyl)-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy]-propyl}-dimethyl-amine;

[3-(5-chloro-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-ylmethoxy)-propyl]-dimethyl-amine;

dimethyl-{3-[3-(1-phenethyl-1H-8-thia-1,3-diaza-dibenzo[e,h]azulen-2-yl)-propoxy]-propyl}-amine; and

a pharmaceutically acceptable salt or solvate thereof.